

Long Day

Film Coated Tablets



Composition:

Each Long Day (30-60) Film Coated Tablet contains: Dapoxetine (as Hydrochloride) 30 mg or 60 mg.

Mechanism of Action:

Dapoxetine is a potent selective serotonin reuptake inhibitor (SSRI), and its major metabolites are equivalent or less potent. The mechanism of action of Dapoxetine in premature ejaculation is presumed to be linked to the inhibition of neuronal reuptake of serotonin and the subsequent potentiation of the neurotransmitter's action at pre- and postsynaptic receptors.

Pharmacokinetics:

Dapoxetine is rapidly absorbed with maximum plasma concentrations (C_{max}) occurring approximately 1-2 hours after tablet intake. The absolute bioavailability is 42%. Dapoxetine and its active metabolite desmethyldapoxetine (DED) are highly bound to plasma proteins. The metabolites of Dapoxetine were primarily eliminated in the urine as conjugates. Following oral administration, the terminal half-life of Dapoxetine and DED are approximately 19 hours.

Indications:

Long Day is indicated for the treatment of premature ejaculation (PE) in adult men aged (18 to 64 years). Long Day should only be prescribed to patients who meet all the following criteria:

- An intravaginal ejaculatory latency time (IELT) of less than two minutes.
- Persistent or recurrent ejaculation with minimal sexual stimulation before, on, or shortly after penetration and before the patient wishes.
- Marked personal distress or interpersonal difficulty as a consequence of Premature ejaculation.
- Poor control over ejaculation and a history of premature ejaculation in the majority of intercourse attempts over the prior 6 months.

Contraindications:

- Hypersensitivity to Dapoxetine or any components of the product.
- Moderate and severe hepatic impairment.
- A history of syncope or a history of mania or severe depression.
- Orthostatic hypotension.
- Long Day is not indicated for use by women.

- Serious heart diseases such as:

- Heart failure (NYHA class II-IV) and significant ischemic heart disease and valvular disease.

- Conduction abnormalities such as atrioventricular block-ing or sick sinus syndrome.

Warnings & Precautions:

- Not intended for the treatment of mental disorders, including depression.
- Caution should be exercised in men with erectile dysfunction who are taking phosphodiesterase-5 inhibitors.
- Not recommended for patients with:
- Controlled epilepsy, a history of bleeding or coagulation disorders.
- Increased intraocular pressure or risk of developing angle-closure glaucoma.
- Severe renal failure (creatinine clearance less than 30 ml/min).
- Mild hepatic impairment (Child-Pugh class A).

Pregnancy & Lactation:

Long Day is not indicated for use by women.

Driving & using machines:

Has minor or moderate influence on the ability to drive and use machines. Caution should be taken when taking Long Day, as it may cause dizziness, fainting, blurred vision, and drowsiness.

Drug Interactions:

- In the case of concomitant use of Long Day with moderate CYP3A4 inhibitors (e.g., clarithromycin, fluconazole, diltiazem); the maximum dose of Dapoxetine should be 30 mg.
- Long Day should be used with caution in patients receiving alpha-adrenergic blockers because of the possibility of orthostatic hypotension.
- The use of Long Day is contraindicated when used concomitantly with:
 - Thioridazine, or within 14 days of stopping thioridazine treatment. Similarly, thioridazine should not be used within 7 days after stopping Long Day.
 - Strong cytochrome CYP3A4 enzymes inhibitors (e.g., ketoconazole, itraconazole, ritonavir, grapefruit juice).
 - Phosphodiesterase-5 inhibitors due to the possibility of orthostatic hypotension.
 - Other SSRIs or MAOIs, or within 14 days of stopping treatment with these preparations. Also, these medicines should not be given within 7 days after stopping Long Day treatment.

Side Effects:

Common side effects are headache, dizziness, anxiety, agitation, restlessness, abnormal dreams, libido decreased, disturbance in attention, tremor, vision blurred, flushing, sinus congestion, diarrhoea, nausea, vomiting, constipation, abdominal pain, dyspepsia, flatulence, dry mouth, erectile dysfunction, excessive sweating and fatigue.

Dosage & Administration:

Adult men (aged 18 to 64 years): The recommended starting dose for all patients is 30 mg, taken as needed approximately 1 to 3 hours prior to sexual activity. Treatment with Long Day should not be initiated with the 60 mg dose.

Long Day is not intended for continuous daily use. Long Day must not be taken more frequently than once every 24 hours.

If the patient's response to 30 mg is insufficient and the patient has not experienced moderate or severe side effects, the dose may be increased to a maximum recommended dose of 60 mg taken as needed approximately 1 to 3 hours prior to sexual activity.

- The safety and effectiveness of Long Day have not been established in patients younger than 18 years of age and Elderly (65 years of age and older).

Overdose:

Symptoms: Somnolence, gastrointestinal disturbances (e.g. nausea, vomiting), tachycardia, dizziness, tremor, and agitation.

Treatment: In cases of overdose, standard supportive measures should be adopted as required. Due to high protein binding and large volume of distribution of Dapoxetine, dialysis is unlikely to be of benefit. No specific antidotes for poisoning Long Day are known.

Storage: Keep out of reach of children. Store below 30°C.

Packaging: Each Long Day (30-60) carton box contains 4 film coated tablets in a blister strip.

